

NEW EFFECTIVE BOUNDS ON THE DIMENSION OF A LINEAR SYSTEM IN $\mathbb{P}^2$

MARCIN DUMNICKI, WITOLD JARNICKI

*Institute of Mathematics, Jagiellonian University,
Reymonta 4, 30-059 Kraków, Poland*

ABSTRACT. The main goal of this paper is to present an algorithm bounding the dimension of a linear system of curves of given degree (or monomial basis) with multiple points in general position. As a result we prove the Hirschowitz–Harbourne Conjecture when the multiplicities of base points are bounded by 11.

Keywords: Multivariate interpolation, Hirschowitz Conjecture.

1. INTRODUCTION

Let $\mathbb{K}$ be a field of characteristic zero, $\mathbb{N} = \{0, 1, 2, \dots\}$, $\mathbb{N}^* = \{1, 2, 3, \dots\}$.

Definition 1. Let $D \subset \mathbb{N}^2$ be finite, let $m_1, \dots, m_r \in \mathbb{N}$, let $p_1, \dots, p_r \in \mathbb{K}^2$. Define the vector space (over $\mathbb{K}$):

$$\mathcal{L}_D(m_1 p_1, \dots, m_r p_r) := \left\{ f = \sum_{\beta \in D} c_\beta X^\beta \mid c_\beta \in \mathbb{K}, \frac{\partial^{|\alpha|} f}{\partial X^\alpha}(p_j) = 0, |\alpha| < m_j, j = 1, \dots, r \right\}.$$

Define the *dimension* of the *system of curves* $\mathcal{L}_D(m_1, \dots, m_r)$ to be

$$\dim \mathcal{L}_D(m_1, \dots, m_r) := \min_{\{p_j\}_{j=1}^r, p_j \in \mathbb{K}^2} \dim_{\mathbb{K}} \mathcal{L}_D(m_1 p_1, \dots, m_r p_r) - 1.$$

Remark 2. If points $p_1, \dots, p_r$ are in general position we have

$$\dim \mathcal{L}_D(m_1, \dots, m_r) = \dim_{\mathbb{K}} \mathcal{L}_D(m_1 p_1, \dots, m_r p_r) - 1.$$

System $\mathcal{L}_D(m_1, \dots, m_r)$ can be understood as a vector space of curves generated by monomials with exponents from D having multiplicities at least $m_1, \dots, m_r$ in r general points.

Let us assume that $D = \{\alpha \in \mathbb{N}^2 \mid |\alpha| \leq d\}$ for some $d \in \mathbb{N}$. Then the space $\mathcal{L}_D(m_1, \dots, m_r)$ can be associated with the linear system over $\mathbb{P}^2$ ($:= \mathbb{P}^2 \mathbb{K}$)

$$\mathcal{L}_d(m_1, \dots, m_r) := |dH| \left(- \sum_{j=1}^r m_j p_j \right).$$

This system contains all divisors from the system $|dH|$ (H being a generic line in $\mathbb{P}^2$) that have multiplicity at least m_j at p_j , where $p_1, \dots, p_r$ are in general position. In particular, the space $\mathcal{L}_D(m_1, \dots, m_r)$ and the linear system $\mathcal{L}_d(m_1, \dots, m_r)$ have equal dimensions.

Intuitively, the dimension of the system $\mathcal{L}_D(m_1, \dots, m_r)$ should be equal to the dimension of $V_D := \text{span}\{X^\alpha \mid \alpha \in D\}$ ($= \#D - 1$) minus the number of conditions imposed by the multiplicities $m_1, \dots, m_r$. However, the actual dimension may be different.

Email address: Witold.Jarnicki@im.uj.edu.pl (corresponding author), Marcin.Dumnicki@im.uj.edu.pl, phone number: +48-12-6339781, fax number: +48-12-6324372.

Definition 3. Let $L = \mathcal{L}_D(m_1, \dots, m_r)$ be a system of curves. Define the *expected dimension* of L

$$\text{edim } L := \max \left\{ \#D - 1 - \sum_{j=1}^r \binom{m_j + 1}{2}, -1 \right\}.$$

Proposition 4. For any system of curves L we have

$$\dim L \geq \text{edim } L.$$

Proof. Fix $p_1, \dots, p_r \in \mathbb{K}^2$. Put

$$\varphi_{j,\alpha}^{p_1, \dots, p_r} : V_D \ni f \mapsto \frac{\partial^{|\alpha|} f}{\partial X^\alpha} (p_j) \in \mathbb{K}, \quad j = 1, \dots, r, \quad \alpha \in \mathbb{N}^2.$$

Let $\mathcal{A} = \{(j, \alpha) \mid |\alpha| < m_j, j = 1, \dots, r\}$. Consider the linear mapping

$$\varphi^{p_1, \dots, p_r} : V_D \ni f \mapsto (\varphi_a^{p_1, \dots, p_r}(f))_{a \in \mathcal{A}} \in \mathbb{K}^{\#\mathcal{A}}.$$

By linear algebra we have $\dim \ker \varphi^{p_1, \dots, p_r} \geq \#D - \#\mathcal{A}$, which implies

$$\dim L = \min_{\{p_j\}_{j=1}^r, p_j \in \mathbb{K}^2} \dim \ker \varphi^{p_1, \dots, p_r} - 1 \geq \text{edim } L.$$

□

Definition 5. We say that *system of curves L is special* if

$$\dim L > \text{edim } L.$$

Otherwise we say that L is *non-special*.

2. THE HIRSCHOWITZ–HARBOURNE CONJECTURE

For systems of the form $\mathcal{L}_d(m_1, \dots, m_r)$ the well-known Hirschowitz–Harbourne Conjecture giving geometrical description to the speciality of a system was formulated in [8]. To formulate this Conjecture consider the blowing-up $\pi : \tilde{\mathbb{P}}^2 \rightarrow \mathbb{P}^2$ in r general points with exceptional divisors $E_1, \dots, E_r$.

Definition 6. A curve $C \subset \mathbb{P}^2$ is said to be *−1-curve* if it is irreducible, and the self-intersection of its proper transform $\tilde{C} \subset \tilde{\mathbb{P}}^2$ is equal to -1 .

Conjecture 7 (Hirschowitz–Harbourne). *A system $L = \mathcal{L}_d(m_1, \dots, m_r)$ is special if and only if there exists a −1-curve $C \subset \mathbb{P}^2$ such that*

$$\tilde{L} \cdot \tilde{C} \leq -2,$$

where $\tilde{L} = |d\pi^*(\mathcal{O}_{\mathbb{P}^2}(1)) - \sum_{j=1}^r m_j E_j|$.

This Conjecture was studied by many authors, we refer only to the recent results. For homogenous systems ($m_1 = m_2 = m_3 = \dots = m_r =: m$), the above conjecture holds for $m \leq 20$ (see [4, 5]). In the general case multiplicities bounded by 7 have been dealt with (see [10]). Both these results were obtained with the help of computers.

For further information about above conjecture see for example [3].

The main result of this paper is Corollary 26 stating that Conjecture 7 holds for multiplicities bounded by 11.

For a given system one can verify whether the above conjecture holds. If a multiple -1 -curve in the base locus cannot be easily found computationally, we check whether the system is non-special by using our reduction method and/or calculating the determinant of φ (see Proposition 4).

Hence, in order to prove Conjecture 7 for linear systems up to multiplicity 11, it is enough to limit the number of cases to be checked. Therefore we will not focus on the conjecture itself, but on the possibility of finding the list of all special systems (in particular we use the reduction method to obtain effective bounds on the number of cases).

3. REDUCTION METHOD


Definition 8. Let $a_1, \dots, a_k \in \mathbb{N}$, $a_j \leq j$, $j = 1, \dots, k$. Define the diagram $(a_1, \dots, a_k)$

$$(a_1, \dots, a_k) = \bigcup_{j=1}^k \{(\alpha_1, \alpha_2) \in \mathbb{N}^2 \mid \alpha_1 + \alpha_2 = j - 1, \alpha_2 < a_j\}.$$

For $a, a_1, \dots, a_k \in \mathbb{N}$, $a_j \leq a + j$ define

$$(\bar{a}, a_1, \dots, a_k) := (1, 2, \dots, a - 1, a, a_1, \dots, a_k).$$

Example 9.



Definition 10. Let $k \in \mathbb{N}^*$, let $D = (b_1, \dots, b_\ell, a_1, \dots, a_k)$ be a diagram, $a_k > 0$. Define the numbers v_j , $j = k, \dots, 1$ inductively to be

$$v_j := \begin{cases} a_j, & a_j < k, \\ \max\{1, \dots, k\} \setminus \{v_{j+1}, \dots, v_k\}, & a_j \geq k. \end{cases}$$

If we have

$$v_i \neq v_j \text{ for } i \neq j, \quad v_j \leq a_j \text{ for } j = 1, \dots, k$$

then we say that D is k -reducible. The diagram

$$\text{red}_k(D) := (b_1, \dots, b_\ell, a_1 - v_1, \dots, a_k - v_k)$$

will be called the k -reduction of D . We use the following notation

$$(b_1, \dots, b_\ell, a_1, \dots, a_k) \xrightarrow{k} (b_1, \dots, b_\ell, a_1 - v_1, \dots, a_k - v_k).$$

Definition 11. Let $k \in \mathbb{N}^*$. We say that a diagram $G = (b_1, \dots, b_\ell, a_1, \dots, a_k)$ is a *weak k -reduction of a diagram $D = (b_1, \dots, b_\ell, a'_1, \dots, a'_k)$* if there exists a sequence of numbers $c_1, \dots, c_k \in \mathbb{N}$ (possibly all of them equal to 0) such that

$$D' := (b_1, \dots, b_\ell, a'_1 + c_1, \dots, a'_k + c_k)$$

is a diagram (in particular $a'_j + c_j \leq \ell + j$, $j = 1, \dots, k$) and $\text{red}_k(D') = G$. We use the following notation

$$(b_1, \dots, b_\ell, a'_1, \dots, a'_k) \xrightarrow{k \text{ w}} (b_1, \dots, b_\ell, a_1, \dots, a_k).$$

Remark 12. Observe that the k -reduction of D is a weak k -reduction.



the 4-reduction of diagram $(\bar{5}, 3, 2)$



a weak 5-reduction of diagram $(\bar{4}, 4, 4)$

Theorem 14. *Let $m_1, \dots, m_r \in \mathbb{N}$. If a diagram D' is a weak m_r -reduction of a diagram D then*

Definition 15. In the situation above, we say that $\mathcal{L}_{D'}(m_1, \dots, m_{r-1})$ is a *weak reduction* of the system $\mathcal{L}_D(m_1, \dots, m_r)$, or simply *the reduction* if $D' = \text{red}_{m_r}(D)$.

Corollary 16. *Let $m_1, \dots, m_r \in \mathbb{N}$. Let $L = \mathcal{L}_D(m_1, \dots, m_r)$. If D is m_r -reducible and the reduction of L is non-special, then L is non-special.*

$$\dim \mathcal{L}_D(m_1, \dots, m_r) \leq \dim \mathcal{L}_G(m_1, \dots, m_r).$$
$$\dim \mathcal{L}_G(m_1, \dots, m_r) \leq \dim \mathcal{L}_{D'}(m_1, \dots, m_{r-1}).$$

Step 1. We will show the following:

- (1) the system $\mathcal{L}_{G \setminus D'}(m_r)$ is non-special,
- (2) if a set $P \subset G$ satisfies $\#P = \#(G \setminus D')$ and

then the system $\mathcal{L}_P(m_r)$ is special.

$$\mathcal{F} = \left\{ F \subset G \mid \#F = \binom{m_r + 1}{2}, F \text{ do not lie on a curve of degree } m_r - 1 \right\}.$$

Step 2. Now we can compute the dimension of a system $L := \mathcal{L}_G(m_1, \dots, m_r)$ as $\dim L = \#G - \text{rank } M - 1$, where M is the matrix of the mapping $\varphi^{p_1, \dots, p_r}$ associated to

L (see the proof of Proposition 4). In an appropriate basis this matrix is of the following form

$$M = \left[\begin{array}{c|c} \begin{array}{c} \text{matrix} \\ \text{associated to} \\ \mathcal{L}_{D'}(m_1 p_1, \dots, m_{r-1} p_{r-1}) \end{array} & K \\ \hline K' & \begin{array}{c} \text{square matrix} \\ \text{associated to} \\ \mathcal{L}_{G \setminus D'}(m_r p_r) \end{array} \end{array} \right].$$

The determinant of the lower right submatrix is nonzero (because of (1)). Take a maximal nonzero minor (of size k) in the upper left submatrix. Then, by generalized Laplace rule and property (2),

$$\text{rank } M = k + \#G - \#D',$$

which finishes the proof. $\square$

In the proof we have used two properties of the set of reduced monomials. In fact these two properties may hold for many other sets, which allows us to find and use other “reductions”.

Definition 17. Let $D \subset \mathbb{N}^2$ be finite (not necessarily a diagram). We say that a system $\mathcal{L}_D(m_1, \dots, m_r)$ admits a reduction algorithm if there exists a sequence of sets $D = D_r \supset \dots \supset D_0$ such that

- (1) $\#D_j - \#D_{j-1} \leq \binom{m_j+1}{2}$, $j = 1, \dots, r$,
- (2) the system $\mathcal{L}_{D_j \setminus D_{j-1}}(m_j)$ is non-special, $j = 1, \dots, r$,
- (3) if $P \subset D_j$ satisfies $\#P = \#D_j - \#D_{j-1}$ and

$$\sum_{\alpha \in P} \alpha = \sum_{\alpha \in D_j \setminus D_{j-1}} \alpha$$

then $\mathcal{L}_P(m_j)$ is special, $j = 1, \dots, r$,

- (4) $\#D_0 - 1 \leq \text{edim } \mathcal{L}_D(m_1, \dots, m_r)$.

The following theorem has been proven in [7] (also for dimension greater than two).

Theorem 18. *If a system admits a reduction algorithm, then it is non-special.*

Moreover, investigating non-speciality of many systems (in dimension two, as well as for higher dimensions) has lead to the following

Conjecture 19. *Every non-special system admits a reduction algorithm.*

Remark 20. In [10] S. Yang presented the “box diagram algorithm”, which can be treated as another possibility of reducing diagrams. However, not every non-special system can be reduced to a trivial one using the reduction of this type. Moreover, Theorem 18 works also in a higher dimension.

Now we can present the main algorithm to bound the dimension of a system.

Algorithm DIMENSIONBOUND

Input: a diagram $D = (\bar{a}, a_1, \dots, a_k)$ and sequence of multiplicities $(m_1, \dots, m_r)$.

Output: a number $d \in \mathbb{N}$ such that $\dim \mathcal{L}_D(m_1, \dots, m_r) \leq d$.

```

for  $i \leftarrow 1$  to  $r$  do {
     $D \leftarrow$  a weak  $m_i$ -reduction of  $D$ 
}

```


$d \leftarrow \#D - 1$
return d

Remark 21. Observe that the quality of the bound obtained depends on the weak reductions chosen. In particular taking D as a weak reduction of D results in a trivial bound on the dimension. Our implementation of this algorithm chooses $D' \supset D$ (see Definition 11) of minimum possible cardinality, which results in much better bounds.

Example 22. Consider the system $L = \mathcal{L}_{27}(24, 6, 6, 6, 6, 6, 6)$. We can see that $\dim L \geq 0$ (picture). The following sequence of weak reductions

$$\begin{aligned} (\overline{28}) &\xrightarrow{24} (\overline{4}, \underbrace{4, \dots, 4}_{24}) \xrightarrow{6 \text{ w}} (\overline{4}, \underbrace{4, \dots, 4}_{18}, 3, 2, 1) \xrightarrow{6 \text{ w}} (\overline{4}, \underbrace{4, \dots, 4}_{15}) \\ &\xrightarrow{6 \text{ w}} (\overline{4}, \underbrace{4, \dots, 4}_{9}, 3, 2, 1) \xrightarrow{6 \text{ w}} (\overline{4}, 4, 4, 4, 4, 4) \xrightarrow{6 \text{ w}} (\overline{4}, 3, 2, 1) \xrightarrow{6 \text{ w}} (1) \end{aligned}$$

gives $\dim L \leq 0$.



The system $L_{27}(24, 6, 6, 6, 6, 6, 6)$. Each curve is triple.

Example 23. We can also use reductions to simplify the computation. One of the cases treated in [4] was the system $L = \mathcal{L}_{133}(20^{\times 43})$. In order to compute the dimension of L using the determinant method, one must consider a 9045×9030 matrix. By reduction, it is enough to compute the dimension of the system $\mathcal{L}_D(20^{\times 5})$, where

$$D = (\overline{36}, 36, 36, 36, 36, 34, 33, 32, 30, 27, 25, 22, 18, 15, 11, 7, 1).$$

Now the matrix is only 1065×1050 , which allows computation of the determinant significantly faster.

4. SPECIAL SYSTEMS WITH BOUNDED MULTIPLICITY

Theorem 24. Let $d, M \in \mathbb{N}^*$, let D be the diagram obtained by applying a sequence of m_j -reductions to the diagram $(\overline{d})$, where $m_j \leq M$. If $\#D > 2M(2M - 1)$ then

- (1) D can be written as $(\overline{a}, a_1, \dots, a_M)$, $a \geq 2M$,
- (2) D is m -reducible for any $m \leq M$.

Proof. For the proof (technical but elementary) see [7]. □

Corollary 25. For a given $m_1, \dots, m_r$ and d large enough, the system $\mathcal{L}_d(m_1, \dots, m_r)$ is non-special.

Proof. Put $M := \max\{m_1, \dots, m_r\}$, take d such that

$$\binom{d+2}{2} > 2M(2M-1) + \sum_{j=1}^r \binom{m_j+1}{2},$$

and apply Theorem 24. $\square$

The above theorem allows us to find all special systems with bounded multiplicities under some conditions. Fix an $M \in \mathbb{N}^*$. The idea is to find a finite set of diagrams $\mathcal{D}$ with the following two properties.

- (1) The system $\mathcal{L}_d(m_1, \dots, m_r)$, where $m_i \leq M$ and d is large enough, reduces to either system $\mathcal{L}_{D'}()$ (without conditions on points) or a system $\mathcal{L}_{D''}(k_1, \dots, k_\ell)$ for $D'' \in \mathcal{D}$,
- (2) Every system $\mathcal{L}_D(k_1, \dots, k_\ell)$, where $D \in \mathcal{D}$ and $k_j \leq M$, $j = 1, \dots, \ell$, is non-special.

The first property can be achieved due to above theorem, which assures that any diagram large enough can be reduced, and describes the form of its reduction. The second property can be checked directly, e.g. by computing determinants and/or reductions.

Having chosen a suitable set $\mathcal{D}$, it is enough to look for special systems in a finite set of systems built on diagrams, which cannot be reduced to one of the diagrams in $\mathcal{D}$.

The effectiveness and low time complexity of computing reductions allowed doing this for multiplicities bounded by 11.

The authors wrote and ran suitable programs to find all special systems up to multiplicity 11. We present here only the numbers of special systems (of respective multiplicities). The list of such systems can be found at [6].

multiplicity	number of special systems
1	0
2	2
3	14
4	91
5	405
6	1798
7	6751
8	25262
9	86147
10	291868
11	929519

As a corollary we have

Corollary 26. *The Harbourne–Hirschowitz Conjecture holds for systems with multiplicities bounded by 11.*

5. BOUNDING THE REGULARITY

Definition 27. The *regularity* of a sequence of multiplicities $(m_1, \dots, m_r)$ is defined to be

$$\text{rg}(m_1, \dots, m_r) := \min \{d \in \mathbb{N} \mid \mathcal{L}_d(m_1, \dots, m_r) \text{ is nonempty and non-special}\}.$$

It can be shown (see Corollary 25) that regularity is a well-defined natural number. We present here a new algorithm to find an upper bound on regularity of a system of multiplicities.

Algorithm REGULARITYBOUND

Input: a sequence of multiplicities $(m_1, \dots, m_r)$.

Output: a number $g \in \mathbb{N}$ such that $\text{rg}(m_1, \dots, m_r) \leq g$.

```

 $g \leftarrow -1$ 
repeat
   $g \leftarrow g + 1$ 
   $d \leftarrow \text{DIMENSIONBOUND}(g, (m_1, \dots, m_r))$ 
until  $d = \text{edim } \mathcal{L}_g(m_1, \dots, m_r) > -1$ 
return  $g$ 

```

Theorem 28. *Algorithm REGULARITYBOUND stops after a finite number of steps and returns the upper bound for the regularity of a given system.*

Proof. The correctness is obvious. The algorithm must stop in view of Corollary 25. $\square$

Definition 29. Let $m_1, \dots, m_r, k_1, \dots, k_r \in \mathbb{N}$. Define

$$(m_1^{\times k_1}, \dots, m_r^{\times k_r}) = (\underbrace{m_1, \dots, m_1}_{k_1}, \dots, \underbrace{m_r, \dots, m_r}_{k_r}).$$

In the paper [9] a new algorithm for bounding the regularity is given. We present the comparison of these two algorithms. We use five sequences of multiplicities (following the author of [9]).

$$\begin{aligned}
L_1 &= (500^{\times 2}, 400, 300, 200^{\times 2}, 150^{\times 3}, 100^{\times 3}, 80^{\times 3}, 10), \\
L_2 &= (350, 300, 250, 200^{\times 2}, 100^{\times 3}, 75^{\times 2}, 70, 60, 50^{\times 2}, 30^{\times 2}), \\
L_3 &= (386, 243, 200, 170^{\times 2}, 162, 100, 81^{\times 3}, 54, 40, 27^{\times 3}, 25, 10, 9^{\times 3}, \\
&\quad 6, 3^{\times 4}, 2, 1^{\times 5}), \\
L_4 &= (600, 350, 300^{\times 3}, 180, 150^{\times 4}, 80^{\times 5}, 50, 40^{\times 4}, 25, 20^{\times 4}, 16, 10^{\times 5}), \\
L_5 &= (500^{\times 3}, 425, 110, 100^{\times 7}, 55, 50^{\times 2}, 45, 20^{\times 5}, 10^{\times 4}).
\end{aligned}$$

For each sequence we give its conjectural regularity (assuming Hirschowitz–Harbourne Conjecture), and the bounds for regularity computed by algorithm of Monserrat, and our reduction method. Moreover, for each system L_i , we present also a pair (d, p) (depending on L_i) such that

$$\dim \mathcal{L}_{(\overline{d+1}, p)}(L_i) = \text{edim } \mathcal{L}_{(\overline{d+1}, p)}(L_i) > -1$$

(also obtained by reduction method).

	L_1	L_2	L_3	L_4	L_5
conjecture	999	650	628	959	1017
[9]	999	676	628	985	1018
reduction	999	650	628	959	1017
reduction	(998, 1)	(649, 100)	(627, 144)	(958, 816)	(1016, 833)

6. REDUCTION AND CREMONA TRANSFORMATION

It is worth mentioning that the Cremona transformation can make reducing much easier.

Definition 30. Let $L = \mathcal{L}_d(m_1, \dots, m_r)$. If L satisfies

$$(1) \ c := m_1 + m_2 + m_3 - d > 0,$$

(2) $m_i \geq c$, for $i = 1, 2, 3$,

then the system $\mathcal{L}_{d-c}(m_1 - c, m_2 - c, m_3 - c, m_4, \dots, m_r)$ is called a *Cremona transformation of L* .

Theorem 31. *For any system $L = \mathcal{L}_d(m_1, \dots, m_r)$, which admits a Cremona transformation we have: L is special if and only if its Cremona transformation is a special system.*

Proof. One can show that if we apply the usual Cremona transformation of $\mathbb{P}^2$ based on points with multiplicities m_1, m_2, m_3 then we obtain a system as above. Of course the dimension cannot change. $\square$

Remark 32. A Cremona transformation does not change the expected dimension of a system (this can be checked by direct computation). The reductions, however, become smaller and easier to perform. Many examples have shown that a combination of Cremona transformation and reduction methods is more effective than using only reductions.

Example 33. Let $L = \mathcal{L}_{63}(24 \times 7)$. If we apply the reduction method to L we obtain that $\dim L \leq 18$. Applying Cremona transformation to L we get the system from example 22, which gives $\dim L = 0$.

REFERENCES

- [1] J. Alexander, A. Hirschowitz, An asymptotic vanishing theorem for generic unions of multiple points, *Invent. Math.* 140, 303–325 (2000)
- [2] J. Alexander, A. Hirschowitz, Polynomial interpolation in several variables, *J. Algebraic Geometry* 4, 201–222 (1995).
- [3] C. Ciliberto, Geometric Aspects of Polynomial Interpolation in More Variables and of Waring’s Problem, *Proc. ECM, Barcelona* (2000).
- [4] C. Ciliberto, F. Cioffi, R. Miranda, F. Orecchia, Bivariate Hermite interpolation and linear systems of plane curves with base fat points, *Computer mathematics, Lecture Notes Ser. Comput.*, 10, World Sci. Publishing, River Edge, NJ, 87–102 (2003).
- [5] C. Ciliberto, R. Miranda, Linear systems of plane curves with base points of equal multiplicity, *Trans. Amer. Math. Soc.* 352, 4037–4050 (2000).
- [6] M. Dumnicki, <http://gamma.im.uj.edu.pl/dumnicki/interpol.htm>, February 10, 2005.
- [7] M. Dumnicki, Expected term bases for generic multivariate Hermite interpolation, *arXiv:math.AG/0503701*.
- [8] A. Hirshowitz, Une conjecture pour la cohomologie des diviseurs sur les surfaces rationelles g n riques, *J. Reine Angew. Math.* 397, 208–213 (1989).
- [9] F. Monserrat, Curves having one place at infinity and linear systems on rational surfaces, preprint.
- [10] S. Yang, Linear systems in $\mathbb{P}^2$ with base points of bounded multiplicity, *arXiv:math.AG/0406591 v2*.